

DYNAMICAL CONSTRAINTS ON EARLY-TYPE GALAXY HALOS

Aaron J. Romanowsky^{1,2}, Magda Arnaboldi³, Nigel G. Douglas¹, Konrad Kuijken^{1,4}, Michael R. Merrifield², Kenneth C. Freeman⁵ and Joris Gerssen⁶

Much insight into the formation of early-type galaxies (ellipticals and S0s) can be gained through study of their halo properties (mass distribution, angular momenta, orbit structure). One of the most promising techniques for probing early-type halos is via the kinematics of planetary nebulae (PNe). We have pursued a multi-pronged program using several different techniques, old and new, to acquire PN velocities. This entails multi-fiber spectroscopy with WYFFOS+AUTOFIB2 on the 4.2-m WHT, masked counter-dispersed imaging with FORS2+MXU on UT2, and counter-dispersed imaging with ISIS and with the custom-built *Planetary Nebula Spectrograph (PN.S)* on the WHT (Douglas et al. 2002). We have now acquired 70-200 velocities to 3-6 R_{eff} in each of 7 galaxies: NGC 821, NGC 3379, NGC 4472, NGC 4486, NGC 4494, NGC 5866, and NGC 7457.

For the bright ellipticals NGC 4472 and NGC 4486, the kinematics of the PNe and the globular clusters (GCs: Zepf et al. 2000; Hanes et al. 2001) suggest that the circular velocity profile is constant or rising with radius at 4 R_{eff} .

For the faint ellipticals NGC 821, NGC 3379, and NGC 4494, the projected velocity dispersion profile declines markedly with radius (see Figure 1). Together with the results for NGC 4697 (Mendéz et al. 2001), this suggests that sub- L^* ellipticals have much less dominant dark halos than brighter ellipticals.

More rigorous determination of the mass distributions will use orbit modeling techniques (Romanowsky & Kochanek 2001) and all available dynamical constraints (stars, PNe, GCs, X-ray gas).

Although elliptical galaxies rotate slowly in their inner parts, it is possible that their outer parts rotate rapidly, making their total specific angular mo-




Fig. 1. Projected velocity dispersion profile around NGC 3379, as traced by the integrated stellar light (*dotted error bars*; Statler & Smecker-Hane 1999) and 110 PNe (*solid error bars*). The solid line shows an isotropic constant mass-to-light ratio Hernquist (1990) model.

menta comparable to those of spiral galaxies (e.g., Arnaboldi et al. 1998). Full analysis of the PN data for our sample galaxies will test this. In NGC 4486, the outer GCs rotate rapidly ($\sim 250 \text{ km s}^{-1}$ for $R > 20 \text{ kpc}$; Côté et al. 2001), while our PN data indicate a lower rotational velocity for the stars ($\sim 100 \text{ km s}^{-1}$). This difference poses a challenge to models of this galaxy's formation.

Observations are planned with the PN.S for 10 more nearby ellipticals to fill out the current sample.

REFERENCES

- Arnaboldi, M., et al. 1998, ApJ, 507, 759
 Côté, et al. 2001, ApJ, 559, 828
 Douglas, N. G., et al. 2002, PASP, submitted
 Hanes, D. A., et al. 2001, ApJ, 559, 812
 Hernquist, L. 1990, ApJ, 356, 359
 Mendéz, R. H., et al. 2001, ApJ, 563, 135
 Romanowsky, A. J., & Kochanek, C. S. 2001, ApJ, 553, 722
 Statler, T. S., & Smecker-Hane, T. 1999, AJ, 117, 839
 Zepf, S. E. et al. 2000, AJ, 120, 2928

¹Kapteyn Astronomical Institute, Postbus 800, 9700 AV Groningen, The Netherlands.

²School of Physics and Astronomy, University of Nottingham, University Park, Nottingham NG7 2RD, England (aaron.romanowsky@nottingham.ac.uk).

³Osservatorio Astronomico di Capodimonte, via Moiariello 16, I-80131 Naples, Italy.

⁴Leiden Observatory, Postbus 9513, 2300 RA Leiden, The Netherlands.

⁵Research School of Astronomy & Astrophysics, MSSSO, ANU, Canberra, Weston Creek, ACT 2611, Australia.

⁶Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, U.S.A.