

Homework Problem #1

1) An $R = 20$ star observed with LRIS (the Keck imaging spectrograph) produces 1890 detected photo-electrons per second. The R-band sky brightness at Mauna Kea is listed at the CFHT WWW site as $20.9 \text{ mag/arcsec}^2$. The LRIS pixel scale is $0.22 \text{ arcseconds/pixel}$, the readout noise is $8e^-$ and the inverse gain of the system is $2.0 e^-/\text{DN}$.

(a) What is the rate of detected e^-/pixel from the sky in the R band?

(b) What is the rate of detected e^- from a $R = 26$ magnitude star observed at an airmass of 1.2 assuming the extinction coefficient in R is $0.1 \text{ mag}/(\text{unit airmass})$?

(c) Assume that you are measuring all of the light for the $R = 26$ magnitude star in an aperture with a radius of 7 pixels. At what exposure time does the measurement become sky dominated?

(d) For the sky-dominated case, how does the S/N scale with exposure time?

(e) How does the S/N scale with seeing (assume you scale the measuring radius linearly with FWHM of point sources).

(f) Make a table of the source noise, sky noise, readnoise and S/N for exposure times of 1, 60, 600, and 3600 seconds.

(g) What is the exposure time required to make an observation of this star with a S/N of 20?

(f) What is the exposure time required to make an observation of this star with $S/N=20$ with WFPC2 in the filter that is the closest match to "R"?