



Hydrogen Burning in More Massive Stars


<http://apod.nasa.gov/apod/astropix.html>




THE CNO CYCLE



Putting it all together, subtracting off the 1.71 MeV carried away by the neutrinos and adding the 2.04 MeV from positron annihilation



The ${}^{12}\text{C}$ is a catalyst. It is not used up but makes the series of reactions possible. Note however, nucleosynthetic aspects.



CNO CYCLE (Shorthand)



nb. $\alpha \equiv {}^4\text{He}$

CNO CYCLE vs PP 1


$$\epsilon_{CNO} \approx 3.4 \times 10^{-4} \rho X_{CNO} X_H (T/10^7)^{20} \text{ erg g}^{-1} \text{ s}^{-1}$$

where X_{CNO} is the mass fraction of carbon, nitrogen, and oxygen combined. This is based on the slowest reaction, $^{14}\text{N}(p,\gamma)^{15}\text{O}$.

$$\begin{aligned} \frac{\epsilon_{CNO}}{\epsilon_{pp}} &= \frac{3.4 \times 10^{-4}}{0.076} \left(\frac{X_{CNO}}{X_H} \right) (T/10^7)^{20-4} \\ &= (4.5 \times 10^{-3} (0.01/0.70)) (T/10^7)^{16} \\ &= 6.4 \times 10^{-5} (T/10^7)^{16} \end{aligned}$$

which is greater than unity for T greater than 18 million K.

This turns out to mean that the CNO cycle dominates in (Population I) stars of over $2 M_{\odot}$.




More Massive Main Sequence Stars

	$10 M_{\odot}$	$25 M_{\odot}$
X_H	0.32	0.35
L	$3.74 \times 10^{37} \text{ erg s}^{-1}$	$4.8 \times 10^{38} \text{ erg s}^{-1}$
T_{eff}	24,800 (B)	36,400 (O)
Age	16 My	4.7 My
T_{center}	$33.3 \times 10^6 \text{ K}$	$38.2 \times 10^6 \text{ K}$
ρ_{center}	8.81 g cm^{-3}	3.67 g cm^{-3}
τ_{MS}	23 My	7.4 My
R	$2.73 \times 10^{11} \text{ cm}$	$6.19 \times 10^{11} \text{ cm}$
P_{center}	$3.13 \times 10^{16} \text{ dyne cm}^{-2}$	$1.92 \times 10^{16} \text{ dyne cm}^{-2}$
% $P_{\text{radiation}}$	10%	33%


Surfaces stable (radiative, not convective); inner roughly 1/3 of mass is convective.

Convective history $15 M_{\odot}$ and $25 M_{\odot}$ stars




The 15 solar mass star is bigger than the 25 solar mass star when it dies

The Sun





<http://www3.kis.uni-freiburg.de/~pnb/granmovtext1.html>



June 5, 1993

Matter rises in the centers of the granules, cools then falls down. Typical granule size is 1300 km. Lifetimes are 8-15 minutes. Horizontal velocities are $1 - 2 \text{ km s}^{-1}$. The movie is 35 minutes in the life of the sun





35 minutes 4680 +/- 50 A filter

size of granules 250 - 2000 km

smallest size set by transmission
through the earth's atmosphere





Image of an active solar region taken on July 24, 2002 near the eastern limb of the Sun.


Looking more towards edge of the sun
to see 3D structure

http://www.boston.com/bigpicture/2008/10/the_sun.html

<http://www.uwgb.edu/dutchs/planets/sun.htm>




sunspots discovered by Galileo and Harriot 1610




A moderately large sunspot. The Earth would just cover the darkest area. The darkest area is called the "umbra". The surrounding radial darkness the "penumbra".

The umbra is cooler by about 1000 K than the surrounding star (note granulation). The magnetic field in the sunspot is typically 1000 - 4000 Gauss. (The Earth's magnetic field is about 1 Gauss; the sun, on the average < 100 Gauss).




Zeeman Effect




Breaks “degeneracy” in energy of states with different “spins”


Strength of the field can be measured by the degree of splitting




1.1 hours



about 25,000 km across




11 year cycle





The sun actually changes its luminosity ...

Solar Cycle Variations



Variability in solar irradiance was undetectable prior to satellite observations starting in 1978

Sunspot activity as a function of latitude




<http://science.mf.nasa.gov/solpad/solar/images/hbf.gif>

NASA/SSSC/HATHAWAY 2004/02

Spots start to happen at high and low latitude and then migrate towards the equator


Radial magnetic field



Hathaway/NASA/MSFC 2009/09

Complete cycle takes 22 years and the sun's magnetic field reverses every 11 years

Little Ice Age



Cause - Multipolar field variation (quadrupole instead of dipole)?
Fluctuations at tacyocline?




The exact cause of the solar cycle is not well understood, but It is known that the magnetic field of the sun (or at least its surface field) goes through reversals every ~11 years. The whole cycle takes 22 years.

In the "Babcock model", the cycle is caused by the *differential rotation of the sun*. In three years the equatorial regions go round 5 additional revolutions compared with the polar ones. This winds up the field and creates stress that is released in part by surface activity (flares, sunspots, etc).


Rotation:
26.8 d at equator
31.8 d at 75° latitude

This differential rotation exists only in the convection zone. The radiative core rotates rigidly.



Babcock Model


Magnetic Field



Babcock Model

Magnetic Field


Differential Rotation



Babcock Model

Magnetic Field


Differential Rotation








Babcock Model

Magnetic Field

Differential Rotation







© Addison Wesley Longman, Inc.




The smallest loop is 3 times the size of the Earth

Energy stored in magnetic field


$$E = B^2 / 8\pi \text{ erg cm}^{-3}$$

Can be released by "reconnection"






http://sprg.ssl.berkeley.edu/~tohban/nuggets/?page=article&article_id=14

Coronal Mass Ejection




Chromosphere



Coronal Loop

Picture at 141 Angstroms

Temperature About 10^6 K




Temperature

Center	15.7 million K
Photosphere	5800 K
Sunspot (umbra)	4240 K
Penumbra	5680 K
Chromosphere	5000 – 20000 K
Corona	0.5 to 3 million K

Density


Mean density entire sun	1.41 g cm ⁻³
Central density	150 g cm ⁻³
Photosphere	10 ⁻⁹ g cm ⁻³
Chromosphere	10 ⁻¹² g cm ⁻³
Corona	10 ⁻¹⁶ g cm ⁻³
Air (earth)	10 ⁻³ g cm ⁻³

Mass loss rate $\sim 10^{-13}$ solar masses/yr



Effects on earth of solar cycle

- Radio communications and satellite health
- Ozone production and hence uv flux at earth
- Cosmic ray flux
- Aurorae



The last reversal was 800,000 yrs ago, but the average time between reversals is 300,000 yrs

Names are the rock strata where the field is measured

<http://www.astronomycafe.net/qadir/q816.html>

The current magnetic field strength at the Earth's surface of 0.6 Gauss. But long term observations show that it is DECREASING at a rate of about 0.07 percent PER YEAR. This means that in 1500 years from now, it will only be about 35 percent as strong as it is today, and in 4000 years it will have a strength of practically zero.



Magnetic north is currently in northern Canada moving at 10 to 50 km/yr.

In a few decades it will reach Siberia


http://science.nasa.gov/science-news/science-at-nasa/2003/29dec_magneticfield/

Solar flares vs solar prominences (the latter are bigger)

<http://science.howstuffworks.com/sun5.htm>

The Solar Neutrino "Problem"

Hydrogen Burning on the Main Sequence



In the sun



- pp1 85%
- pp2 15%
- pp3 0.02%

$T_{\text{central}} = 15.7$ Million K


Hydrogen Burning on the Main Sequence

In all cases $4p \rightarrow {}^4\text{He} + 2\nu_e + 2e^+$



pp1



pp2




pp3



Neutrino Energies

Species	Average energy	Maximum energy	
p+p	0.267 MeV	0.420 MeV	
${}^7\text{Be}$	0.383 MeV 0.861	0.383 MeV 0.861	10% 90%
${}^8\text{B}$	6.735 MeV	15 MeV	

In the case of ${}^8\text{B}$ and p+p, the energy is shared with a positron hence there is a spread. For ${}^7\text{Be}$ the electron capture goes to a particular state in ${}^7\text{Li}$ and the neutrino has only one energy




Since 1965, experiments have operated to search for and study the neutrinos produced by the sun - in order to:


- Test solar models
- Determine the central temperature of the sun
- The flux of neutrinos from ${}^8\text{B}$ is sensitive to T^{18}
- Learn new particle physics

DETECTORS


The chlorine experiment – Ray Davis – 1965 - ~1999



The gallium experiments (GALLEX and SAGE) –
1991 – 1997 and 1990 – 2001



Kamiokande II - 1996 – 2001



Inelastic scattering of neutrinos on electrons in water. Threshold 9 MeV. Scattered electron emits characteristic radiation.



Homestake Gold Mine
Lead, South Dakota

4850 feet down

tank 20 x 48 feet
615 tons (3.8 x 10⁵ liters)
C₂Cl₄


Threshold 0.814 MeV

Half-life ³⁷Ar = 35.0 days

Neutrino sensitivity
⁷Be, ⁸B

8 x 10³⁰ atoms of Cl


Nobel Prize 2002



In Gran Sasso Tunnel – Italy

3300 m water equivalent

30.3 tons of gallium in GaCl₃-
HCl solution




Threshold 0.233 MeV



Sees pp, ⁷Be, and ⁸B.

Calibrated using radioactive ⁵¹Cr neutrino source


Kamiokande II (in Japanese Alps) 1996 - 2001



Depth 1 km
Detector H₂O
Threshold 9 MeV
Sensitive to ⁸B
20' photomultiplier
tubes
Measure Cerenkov
light
2.3 x 10³² electrons




The Sun - 1999
 (First picture in neutrinos)



This "picture" was taken using data from the Kamiokande 2 neutrino observatory. It contains data from 504 nights (and days) of observation. The observatory is about a mile underground.

Each pixel is about a degree and the whole frame is 90° x 90°.

Sudbury Neutrino Observatory



6800 ft down
 1000 tons D₂O.
 20 m diameter
 Sudbury, Canada
 Threshold 5 MeV

Sees ⁸B decay but can see all three kinds of neutrinos


ν_e, ν_μ, ν_τ

Particle physics aside:

Three Generations of Matter (Fermions)				
	I	II	III	
mass	2.4 MeV	1.27 GeV	171.2 GeV	0
charge	$\frac{2}{3}$	$\frac{2}{3}$	$\frac{2}{3}$	0
spin	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
name	u up	c charm	t top	γ photon
	4.8 MeV	104 MeV	4.2 GeV	0
	$-\frac{1}{3}$	$-\frac{1}{3}$	$-\frac{1}{3}$	0
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
Quarks	d down	s strange	b bottom	g gluon
	<2.2 eV	<0.17 MeV	<15.5 MeV	91.2 GeV
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	0
emitted by pp-cycle cosmology limits the sum of the 3 neutrino masses to < 1 eV	ν_e electron neutrino	ν_μ muon neutrino	ν_τ tau neutrino	Z⁰ weak force
	0.511 MeV	105.7 MeV	1.777 GeV	80.4 GeV
	-1	-1	-1	±1
Leptons	e electron	μ muon	τ tau	W[±] weak force

Bosons (Forces)


Total Rates: Standard Model vs. Experiment
Bahcall-Pinsonneault 2000




<http://www.sno.phy.queensu.ca/sno/sno2.html> - interactions

Neutrino interactions with heavy water D₂O = ²H₂O


Electron neutrino



All neutrinos



add salt to increase sensitivity to neutrons,



Results from SNO – 2002

The flux of electron flavored neutrinos above 5 MeV (i.e., only pp3 = ⁸B neutrinos) is

$$1.76 \pm 0.1 \times 10^6 \text{ cm}^{-2} \text{ s}^{-1}$$

But the flux of μ and τ flavored neutrinos is

$$3.41 \pm 0.64 \times 10^6 \text{ cm}^{-2} \text{ s}^{-1}$$

Nobel Prize in Physics - 2002

Standard Solar Model ⁸B neutrinos

$$5.05^{+1.01}_{-0.81} \times 10^6 \text{ neutrinos cm}^{-2} \text{ s}^{-1}$$

The explanation of the solar neutrino “problem” is apparently *neutrino flavor mixing*.

A flux that starts out as pure electron-”flavored” neutrinos at the middle of the sun ends up at the earth as a mixture of electron, muon, and tauon flavored neutrinos in comparable proportions.

The transformation occurs in the sun and is complete by the time the neutrinos leave the surface. The transformation affects the highest energy neutrinos the most (MSW-mixing).

Such mixing requires that the neutrino have a very small but non-zero rest mass. This is different than in the so called “standard model” where the neutrino is massless. The mass is less than about 10^{-5} times that of the electron.

New physics... (plus we measure the central temperature of the sun very accurately – 15.71 million K)