

## *Astronomy 12*

### *Stars, Stellar Evolution, and High Energy Astrophysics*

<http://www.ucolick.org/~woosley/>  
<http://apod.nasa.gov/apod/astropix.html>

- Stan Woosley

## *The Nature of Astronomy*

- *The scientific study of objects beyond earth (here with emphasis on stars and physics)*
- *A progress report. Our views of the cosmos change daily (but the new theories often include the old ones as subsets)*
- *The cosmos itself changes; all of its constituents are evolving.*
- *A novel aspect of astronomy is its ability to carry out direct studies of the past*
- *Interesting experiments are set up for us, but we have no control over them. Everyone is an observer!*

See also Fraknoi, Morrison and Wolff  
*Prologue*

*The scientific mind does not so much  
provide the right answers as ask the  
right questions*

Claude Levi-Strauss  
French philosopher  
1908 - 2009

*One thing I have learned in a long life is  
that all our science, measured against reality,  
is childlike - and yet it is the most precious  
thing we have.*

Albert Einstein  
Physicist  
1879 - 1955

## *“Astrophysics”*

*The universe obeys physical laws and  
those laws do not vary with space or time*

*It is best understood on the basis of  
physical “models” and mathematics*

### Scientific notation

$$\begin{array}{ll} 1 = 1.0 \times 10^0 & 0.10 = 1.0 \times 10^{-1} \\ 10 = 1.0 \times 10^1 & 0.00000010 = 1.0 \times 10^{-7} \\ 1,000,000 = 1.0 \times 10^6 & 0.00346 = 3.46 \times 10^{-3} \\ 3,450,000 = 3.45 \times 10^6 & 0.002356347 \approx 2.36 \times 10^{-3} \end{array}$$

$$(1.0 \times 10^{-2})(2.0 \times 10^4) = 2.0 \times 10^2 = 200$$

$$(1.0 \times 10^{-2}) / (2.0 \times 10^4) = \frac{1}{2} \times 10^{-6} = 5.0 \times 10^{-7}$$

In Ay12 (e.g. homework), use only the precision justified by the statement of the problem. The default is 3 figures of accuracy.

### Logarithms

$$\begin{array}{ll} \log(1) = 0 & \log(10) = 1 \\ \log(0.1) = -1 & \log(10^{-6}) = -6 \\ \log(52.3) = 1.72... & \\ \log(a)(b) = \log(a) + \log(b) & \log(a^b) = b \log(a) \\ \text{e.g. } \log(10^2) = 2 \log(10) = 2 & \\ \log(10^x) = x \log(10) = x & \\ \log(100) = \log(10)(10) = \log(10) + \log(10) = 2 & \end{array}$$

logarithms are used extensively in the stellar magnitude system because of the need to describe brightnesses that span many orders of magnitude.

### Angular Measure

(used, e.g., for distance determination)

$$\begin{array}{l} 1 \text{ full circle} = 360 \text{ degrees} \\ 1 \text{ degree} = 60 \text{ arc minutes} \\ 1 \text{ arc minute} = 60 \text{ arc seconds} \end{array}$$

[http://mintaka.sdsu.edu/GF/explain/atmos\\_refr/angles.html](http://mintaka.sdsu.edu/GF/explain/atmos_refr/angles.html)

$$2\pi \text{ radians} = 360 \text{ degrees}$$


A *radian* is the angle subtended by a length of arc equal to the radius of the circle

$$1 \text{ radian} = 360/2\pi = 57.29... \text{ degrees} = 206,265 \text{ arc seconds}$$

Length of arc,  $s$ , subtended by angle  $\theta$

$$s = r \theta$$

if  $\theta$  is measured in radians



Thumb at arm's length  $\sim 2$  degrees

Little finger at arm's length  $\sim 1$  degree

hand spread  $\sim 20$  degrees

Smallest angle with naked eye  $\sim 1$  arc min

Sun or moon  $\sim 1/2$  degree

HST  $\sim 0.4$  milli-arc-seconds (0.01 pixel – for astrometry)

GAIA  $\sim 10 - 20 \mu\text{as}$  (planned, by mission's end; 1 foot at 2 million miles)

### Units

The basic units in Ay12 are cm, gm, and sec (with apologies to the physicists).

How many cm in a light year?  $1 \text{ ly} = c * 1 \text{ yr}$


$c = 2.99 \times 10^8 \text{ km s}^{-1}$  based on Julian year = 365.25 days (exactly)  
day = 24 hours

km =  $10^3$  m hour = 60 minutes  
m =  $10^2$  cm

$1 \text{ ly} = c \times 1 \text{ year}$

$$\approx \left( \frac{2.99 \times 10^8 \text{ km}}{\text{s}} \right) (1 \text{ yr}) \left( \frac{10^3 \text{ m}}{\text{km}} \right) \left( \frac{10^2 \text{ cm}}{\text{m}} \right) \left( \frac{365 \text{ day}}{1 \text{ yr}} \right) \left( \frac{24 \text{ hr}}{1 \text{ day}} \right) \left( \frac{60 \text{ min}}{1 \text{ hr}} \right) \left( \frac{60 \text{ s}}{1 \text{ min}} \right)$$
$$\approx 9.44 \times 10^{17} \text{ cm}$$

### Spherical Geometry



Circumference of a circle =  $2\pi r$

Surface area of a sphere =  $4\pi r^2$

Volume of a sphere =  $\frac{4}{3}\pi r^3$

Mass of a sphere with *constant density*  $\rho$

$$M = \left( \frac{4}{3}\pi r^3 \right) \rho$$

*To a good approximation stars are spheres*

Mass of a sphere with radius  $r$  and constant density  $\rho$  ( $\text{gm cm}^{-3}$ )

$$M = \frac{4}{3}\pi r^3 \rho$$

E.g., How much does a (spherical) asteroid with radius 5 km and density  $5 \text{ gm cm}^{-3}$  “weigh”?

$$M = \frac{4}{3}(3.14)(5 \times 10^5 \text{ cm})^3 (5 \frac{\text{gm}}{\text{cm}^3})$$
$$= (4.19)(125 \times 10^{15})(5) \text{ gm}$$
$$= 2.62 \times 10^{18} \text{ gm}$$

### Calculus


$$\frac{d}{dx} x^n = n x^{n-1} \quad \int x^n dx = \left( \frac{x^{n+1}}{n+1} \right)$$

$$\frac{d}{d\theta} (\text{Cos } \theta) = -\text{Sin } \theta \quad \frac{d}{d\theta} (\text{Sin } \theta) = \text{Cos } \theta$$

Binomial expansion theorem

$$(1 + \epsilon)^n \approx (1 + n\epsilon) \quad \text{if } |\epsilon| \ll 1$$

e.g.  $(1 + 0.01)^{\frac{1}{2}} \approx 1.005$  (actually 1.0049876..)



Eg. Volume of a sphere

Area of a shell =  $4 \pi r^2$

Thickness =  $\Delta r \approx dr$

Add up a whole bunch of shells

$$\int_0^{r_0} 4\pi r^2 dr = \frac{4}{3}\pi r_0^3$$

#### ANGULAR MEASURE

$\pi = 3.14159\dots$   
 $2 \pi$  radians =  $360^\circ$   
 1 radian =  $57^\circ.296$   
 1 degree =  $60'$  = 60 arc min  
 1 arc min =  $60''$  = 60 arc sec  
 1 radian =  $206265''.806$   
 Number of square degrees on sky =  $41,252,961$

#### PHYSICAL CONSTANTS

Speed of light	$c$	$2.99792 \times 10^{10}$ cm s <sup>-1</sup>
Constant of gravitation	$G$	$6.672 \times 10^{-8}$ dyne cm <sup>2</sup> g <sup>-2</sup>
Planck's constant	$h$	$6.626 \times 10^{-27}$ erg s
Boltzmann's constant	$k$	$1.381 \times 10^{-16}$ erg (deg K) <sup>-1</sup>
Mass hydrogen atom	$m_H$	$1.673 \times 10^{-24}$ g
Avogadro's number	$N_A$	$6.022 \times 10^{23}$ g <sup>-1</sup>
Mass electron	$m_e$	$9.1095 \times 10^{-28}$ g
Charge on the electron	$e$	$4.803 \times 10^{-10}$ electrostatic units
Stefan-Boltzmann radiation constant	$\sigma$	$5.670 \times 10^{-5}$ erg cm <sup>-2</sup> s <sup>-1</sup> (deg K) <sup>-4</sup>
Radiation energy density constant	$a = 4\sigma/c$	$7.56 \times 10^{-15}$ erg cm <sup>-3</sup> (deg K) <sup>-4</sup>
Constant in Wien's Law	$\lambda_{max} T$	$0.28979$ cm (deg K) <sup>-1</sup>
Electron volt	eV	$1.6022 \times 10^{-12}$ erg
Million electron volts	MeV	$10^6$ eV
Angstrom	Å	$10^{-8}$ cm
1 Megaton of TNT	MT	$4.2 \times 10^{22}$ erg

#### ASTRONOMICAL CONSTANTS

Astronomical Unit	AU	$1.495978707 \times 10^{13}$ cm
Parsec	pc	206265 AU
		3.262 ly
		$3.086 \times 10^{28}$ cm
Light year	ly	$9.4605 \times 10^{17}$ cm
		$6.324 \times 10^4$ AU
(sidereal) year	yr	$3.155815 \times 10^7$ s
Mass of Earth	$M_E$	$5.977 \times 10^{27}$ g
(Equatorial) radius of Earth	$R_E$	$6.378 \times 10^8$ cm
Mass of sun	$M_\odot$	$1.989 \times 10^{33}$ g
Radius of sun	$R_\odot$	$6.960 \times 10^{10}$ cm
Luminosity of sun	$L_\odot$	$3.83 \times 10^{33}$ erg s <sup>-1</sup>
Solar constant at Earth	$S$	$1.37 \times 10^8$ erg cm <sup>-2</sup> s <sup>-1</sup>

#### MISCELLANEOUS

Area circle	$A = \pi R^2$
Area of a sphere	$A = 4\pi R^2$
Volume of a sphere	$V = \frac{4}{3}\pi R^3$
Latitude of Santa Cruz	36.9998 degrees N
Longitude of Santa Cruz	122.0624 degrees W
Temperature in K	Temperature in C + 273.15
Temperature in F	(Temperature in C) * 9/5 + 32
5 magnitudes	factor of 100 in flux
For very small $\theta \ll 1$ radian	$\sin \theta \approx \tan \theta \approx \theta$

*Our location in the  
Universe*

### “Spaceship Earth”



From Apollo 11  
1969

### The Earth as a planet

- $M_{\text{earth}} = 5.997 \times 10^{27} \text{ gm}$
- $R_{\text{earth}} = 6.378 \times 10^8 \text{ cm}$
- Age  $\sim 4.54$  billion years (U,Th dating - close to age of sun)
- Orbit sun =  $1.496 \times 10^{13} \text{ cm}$  ( $\sim$ average distance) = **AU**  
(93 million miles) [prior to 1976 was semi-major axis; now radius of circular orbit with the equivalent period]
- Period around the sun = 365.242199... days  
(Julian year = 365.25 days; 86,400 s; exactly)
- Average density =  $5.52 \text{ gm/cm}^3$   $\rho \approx \left( \frac{M}{4/3\pi r^3} \right)$   
densest planet in the solar system,  
barely beats Mercury

aside – leap year

Every year that is divisible by 4

but now years divisible by 100 unless they  
are divisible by 400

e.g., 2100 will not be a leap year  
2000 was a leap year


After 8000 years this system will be off by a day



*or a big, rusty, sandy rock....*

**34.6% Fe**  
**29.5% O**  
**15.2% Si**  
**12.7% Mg**  
**2.4% Ni**  
**1.9% S**

In contrast to .....



*Where did these elements come from?*

*This rusty sandy rock orbits  
the nearest star, the sun....*

### The Sun

*The only star we can study in great detail*


Mass =  $1.989 \times 10^{33}$  gm; about 300,000 Earth masses  
Radius =  $6.96 \times 10^5$  km; almost 100 Earth radii  
Average density  $1.41 \text{ gm/cm}^3$   
Age =  $4.567 \times 10^9$  years  
Luminosity =  $3.90 \times 10^{33}$  erg/s  
(world's armament in  $10^{-5}$  seconds)  $1.37 \times 10^6 \text{ erg cm}^{-2} \text{ s}^{-1}$   
at earth

Central temperature = 15.7 million K  
Photospheric temperature about 5700 K  $K = C + 273$

Rotation period 24.47 days at the equator  
slower near poles

Surface composition (by mass) 70.6% H  
27.5% He, 1.9% C, N, O, Fe, Si, etc  
(like "universe")


**A typical star. A little on the heavy side.**



The figure at the left shows the effect of zooming out in distance from our solar system by a total factor of 100,000 ( $10^5$ ).

At this scale the next star system over, alpha-centauri, becomes visible.

Most of the universe, even within galaxies, is empty.



The nearest (24) stars within 12.5 light years of the earth. The closest star system – Alpha Centauri – is about 7000 times the radius of Pluto’s orbit. 270,000 times the radius of the Earth’s orbit,

circles indicate plane of Milky Way galaxy


Some specific nearby stars:

- *The sun* – a typical yellow dwarf star. Type G2 with 8 planets
- *Proxima Centauri* – closest of the triplet of stars loosely known as “alpha-Centauri”  
Proxima Centauri is a faint red star that orbits Alpha-Centauri A and B with a period of about one million years. Proxima Centauri is 4.22 light years from the Earth (now) and about 0.24 light years from Alpha-Centauri A and B.
- *Alpha-Centauri A and B* – a double star system with a period of about 80 years. Component A is a near twin of the sun (Type G2). Component B is a little fainter and orange. Alpha-Centauri A and B are 4.39 light years from the Earth.
- *Barnards star* – highest proper motion of all stars. 5.9 light years away. It moves 0.29 degrees per century. In another 8000 years Barnard’s star will be the closest star to us (3.8 ly in 11700 AD). M star, faint, red, about 11 Gyr old. No big planets.
- *Lalande 21185* – One of the brightest red dwarfs in the sky but still requires binoculars to see it. In 1996 a couple of Jupiter sized planets were discovered here
- *Epsilon Eridani* – 10.5 light years away. Searched for life by radio searches in the 1960’s. May have a Jupiter sized planet orbiting at a distance of 3.2 AU. Young star (1Gyr?). K2
- *Procyon A,B* – 11.41 light years away. Another multiple star system. 8<sup>th</sup> brightest star in the sky has a white dwarf companion
- *Sirius A,B* – At a distance of 8.60 light years Sirius A is the brightest star in the sky. Sirius B is a white dwarf

Brightest stars		Nearest Stars	
Star name	Apparent magnitude	Star name	distance (ly)
Sun	-26.8	Sun	-
Sirius	-1.46	Proxima Centauri	4.2
Canopus	-0.72	Alpha Centauri AB	4.3
Arcturus	-0.04	Barnards stars	6.0
Alpha Centauri	-0.01	Wolf 359	7.7
Vega	0.00	BD 36+2147	8.2
Capella	0.08	Luyten 726-8AB	8.4
Rigel	0.12	Sirius A B	8.6
Procyon	0.38	Ross 154	9.4
Betelgeuse	0.41	Ross 248	10.4

most nearby stars are too faint to see without a telescope


*In a volume limited sample – counting all stars...*



*Most stars are less luminous than the sun, only a few are brighter.*

### Masses and luminosities

*In binary star systems we can determine the mass of the star. For stars that are spectroscopically "main sequence" the star's luminosity is correlated with its mass.*



*e.g., 10 solar masses is between  $10^3$  and  $10^4$  times more luminous than the sun. 0.1 solar masses is down by  $10^3$*


To summarize:

- There are many more faint stars than bright ones
- Faint stars also have low mass
- Low mass stars live a long time

The converse is also true:

- Bright (high luminosity) stars are rare
- Bright stars are more massive (exception red giants)
- Massive stars have short lives

20 light years – 78 systems – 109 stars




isotropic distribution

50 ly  
~2000 stars



250 light years



Starting to see some preference for Galactic plane for distances beyond this.

Number for isotropic distribution and constant density  
 $n \propto d^3$


About 250,000 stars lie within 250 light years of the Earth. Beyond this distance it becomes difficult to see all the stars in the plane of the Milky Way Galaxy because of the presence of dust.

Only the 1500 most luminous of these stars are plotted. Most of these are visible to the unaided eye.

Note the presence of the Hyades cluster.

< 1500 stars are visible to the unaided eye. More often it's a few hundred.


The Hyades Open cluster of stars (151 light years)



The bright red star Aldebaran is not in the Hyades


This cluster of stars is only about 625 million years old and is in the process of coming apart. Stars like this are born together from a giant cloud of molecular gas, most of which is blown away by the young stars. About 200 stars are catalogued at [http://en.wikipedia.org/wiki/List\\_of\\_stars\\_in\\_Hyades](http://en.wikipedia.org/wiki/List_of_stars_in_Hyades)

5000 light year view – Galactic spiral arm structure is becoming apparent. The sun is on the “Orion Arm” a lesser arm of the Milky Way compared e.g., to the Sagittarius Arm. There is also a lot of gas and dust.



Betelgeuse 650 ly; Orion 1350 ly

The entire visible galaxy is about 80,000 light years across. Note orbiting galaxy and globular clusters



<http://www.atlasoftheuniverse.com/galaxy.html>

### Globular Clusters




47 Tuc

Second brightest globular cluster (behind Omega Cen). There are about 200 globular clusters altogether. This one is near the direction of the SMC in the sky and about 20,000 ly distant. Lots of red giants visible here.

M13

This globular cluster in Hercules is 22,000 ly distant and contains  $10^5 - 10^6$  stars. Age ~ 12 to 14 billion years. It is about 150 light years across.

The clearest experimental evidence for spiral structure in our own galaxy comes from radio observations. The galaxy is transparent in the 21 cm line of atomic H.




the sun

### Radio View of the Milky Way


Interstellar dust does not absorb radio waves

We can observe any direction throughout the Milky Way at radio waves




Radio map at a wavelength of 21 cm, tracing neutral hydrogen

<http://www.geekosystem.com/wise-all-sky-atlas/>



Released 2012. Wide Field Infra-Red Survey Explorer (WISE) composite photograph of the entire sky. Over 500 million individual stars catalogued, though not with great precision.



Hipparcos Space Astrometry Mission (1989 – 1993)

Catalogue of accurate distances (1 milli arc s angular resolution)

118,218 stars


Total stars observed (Tycho 2 Catalogue; 25 mas)

2,539,913 stars

Including 99% of all stars brighter than 11<sup>th</sup> magnitude

*Aside: In AD 150 Ptolemy in his Almagest catalogued 1,022 of the brightest stars.*


Hipparcos – number of stars per square degree (there are 41,253 square degrees in the sky)



The center of our galaxy is towards the constellation Sagittarius




Infrared observation (2MASS) towards center of the Milky Way - dust glows in IR



Optical - M 6,7,18,21,23,24,24 = open clusters; M 16,17,20 = nebulae; M 9, 22,28,54,69,70 = globular clusters

GAIA – just launched (12/19/13) will study the properties of *one billion* stars. Each will be observed 70 times during its 5 year mission. Will obtain accurate distances for 100,000 stars, less accurate distances for one million





Other *spiral* galaxies are thought to look very similar to our own Milky Way.



## History of the Milky Way




The traditional theory:

Quasi-spherical gas cloud fragments into smaller pieces, forming the first, metal-poor stars (pop. II);

Rotating cloud collapses into a disk-like structure

Later populations of stars (pop. I) are restricted to the disk of the Galaxy

## Changes to the Traditional Theory




Ages of stellar “populations” may pose a problem to the traditional theory of the history of the Milky Way

Possible solution: Later accumulation of gas, possibly due to mergers with smaller galaxies

Recently discovered ring of stars around the Milky Way may be the remnant of such a merger

<http://www.sdss.org/news/releases/20030106.milkyway.html>

Sloan Digital Sky Survey



*About 3 billion years in the future, our galaxy and Andromeda will merge.*

*Calculation by John Dubinsky at CITA.*

*Galaxies collide ....*



*The Antenna Galaxy is not one but two galaxies in the process of merging.*

*New generations of stars are being born, even new globular clusters, in the blue regions. Note also the presence of a lot of dust.*

Besides spiral galaxies like Andromeda ...  
(2.2 Mly)



Similar to but somewhat larger and brighter than the Milky Way (has about 250 globular clusters and many orbiting dwarf ellipticals)

There are also Elliptical galaxies



For example, the massive elliptical galaxy M87 at the center of the Virgo cluster of galaxies.

Such galaxies are oval in shape, have no discernible spiral structure, and little gas or dust.

Reddish in color. Very few new stars being born.

Elliptical galaxies come in all sizes from just a little larger than globular clusters to 10 times the mass of the Milky Way.

The most common kind of galaxy nowadays are the dwarf ellipticals.

Gas used up long ago making stars or stripped by galactic collisions and encounters.

## Irregular and other galaxies

The SMC contains several hundred million stars



Small Magellanic Cloud  
AAG ref URS 17



NGC 3102  
AAG ref AAT 41



NGC 1143 irregular galaxy  
AAG ref AAT 46



Large Magellanic Cloud  
AAG ref URS 14




NGC 6022  
AAG ref AAT 28

The LMC (157,000 ly) is the fourth largest galaxy in the local group and contains about 10 billion solar masses

- 1) Andromeda
- 2) Milky Way
- 3) Triangulum Galaxy (M33)
- 4) LMC




## The nearest members of the Local Group of Galaxies orbit our Milky Way




(500,000 ly)

### Fornax dwarf galaxy

460,000 ly distant, discovered in 1938




Like most dwarf galaxies it doesn't look very impressive. Contains only a few million stars. Orbits by six globular clusters



### Clusters of Galaxies


#### Rich clusters of galaxies

- thousands of galaxies
- concentrated toward the center
- more ellipticals
- hot gas
- lots of mergers




#### Poor clusters of galaxies

- just a few galaxies
- ragged shapes
- more spirals,
- fewer ellipticals



### One Billion Light Years




7% of the radius of the visible universe

- 80 superclusters
- 160,000 galaxy groups
- 3 million large galaxies
- 30 million dwarf galaxies
- 500 million billion stars

The nearest really large supercluster is in Centaurus. Virgo is small by comparison.

Structure starting to be filamentary with walls and voids.

### 2dF Galaxy Redshift Survey - AAT (2003) closest 3 billion light years



Survey obtained spectra for 232,155 galaxies over 272 nights of observation. 1500 square degrees. Did not sample the whole sky. On the whole the universe is *homogeneous and isotropic*. Note cellular structure.

<http://www.mso.anu.edu.au/2dFGRS/>



*The Hubble eXtreme Deep Field photo made from a composite of 2000 images taken over a decade. Like a 23 day time exposure*

*Shows a piece of the universe when it was "only" 400 million years old. 5500 galaxies*


*The current age is 13.8 billion years*

### Hubble eXtreme Deep Field 9/2012 $z = 10.3$






<http://www.mpa-garching.mpg.de/galform/virgo/millennium/>



Numerical simulation of cosmic structure  
"The millenium simulation project"

<http://www.mpa-garching.mpg.de/galform/virgo/millennium/>

Approximately 15 Billion Light Years (a sketch)



~300,000 superclusters

~ $10^{10}$  large galaxies

~ 2000 billion billion stars

*The end of the road  
(for now) ...*