



Hydrogen Burning in More Massive Stars and The Sun


<http://apod.nasa.gov/apod/astropix.html>




THE CNO CYCLE




Putting it all together, subtracting off the 1.71 MeV carried away by the neutrinos and adding the 2.04 MeV from positron annihilation



The ^{12}C is a catalyst. It is not used up but makes the series of reactions possible. Note however, nucleosynthetic aspects.



CNO CYCLE (Shorthand)



nb. $\alpha \equiv ^4\text{He}$

CNO CYCLE vs PP 1


$$\epsilon_{CNO} \approx 3.4 \times 10^{-4} \rho X_{CNO} X_H (T/10^7)^{20} \text{ erg g}^{-1} \text{ s}^{-1}$$

where X_{CNO} is the mass fraction of carbon, nitrogen, and oxygen combined. This is based on the slowest reaction, $^{14}\text{N}(p,\gamma)^{15}\text{O}$.

$$\begin{aligned} \frac{\epsilon_{CNO}}{\epsilon_{pp}} &= \frac{3.4 \times 10^{-4}}{0.076} \left(\frac{X_{CNO}}{X_H} \right) (T/10^7)^{20-4} \\ &= (4.5 \times 10^{-3} (0.01/0.70)) (T/10^7)^{16} \\ &= 6.4 \times 10^{-5} (T/10^7)^{16} \end{aligned}$$

which is greater than unity for T greater than 18 million K.


This turns out to mean that the CNO cycle dominates in (Population I) stars of over $2 M_{\odot}$.




More Massive Main Sequence Stars

	$10 M_{\odot}$	$25 M_{\odot}$
X_H	0.32	0.35
L	$3.74 \times 10^{37} \text{ erg s}^{-1}$	$4.8 \times 10^{38} \text{ erg s}^{-1}$
T_{eff}	24,800 (B)	36,400 (O)
Age	16 My	4.7 My
T_{center}	$33.3 \times 10^6 \text{ K}$	$38.2 \times 10^6 \text{ K}$
ρ_{center}	8.81 g cm^{-3}	3.67 g cm^{-3}
τ_{MS}	23 My	7.4 My
R	$2.73 \times 10^{11} \text{ cm}$	$6.19 \times 10^{11} \text{ cm}$
P_{center}	$3.13 \times 10^{16} \text{ dyne cm}^{-2}$	$1.92 \times 10^{16} \text{ dyne cm}^{-2}$
% $P_{\text{radiation}}$	10%	33%

Surfaces stable (radiative, not convective); inner roughly 1/3 of mass is convective.




The Sun




Convection zone ~2% of mass; ~25% of radius

<http://www3.kis.uni-freiburg.de/~pnb/granmovtext1.html>




June 5, 1993

Matter rises in the centers of the granules, cools then falls down. Typical granule size is 1300 km. Lifetimes are 8-15 minutes. Horizontal velocities are $1 - 2 \text{ km s}^{-1}$. The movie is 35 minutes in the life of the sun



Andrea Malagoli



35 minutes 4680 +/- 50 A filter

size of granules 250 - 2000 km

smallest size set by transmission
through the earth's atmosphere






Image of an active solar region taken on July 24, 2002 near the eastern limb of the Sun.

http://www.boston.com/bigpicture/2008/10/the_sun.html

<http://www.uwgb.edu/dutchs/planets/sun.htm>





sunspots discovered by Galileo and Harriot 1610




A moderately large sunspot. The Earth would just cover the darkest area. The darkest area is called the "umbra". The surrounding radial darkness the "penumbra".

The umbra is cooler by about 1000 K than the surrounding star (note granulation). The magnetic field in the sunspot is typically 1000 - 4000 Gauss. (The Earth's magnetic field is about 1 Gauss; the sun, on the average < 100 Gauss).




2 Detailed closeup of magnetic structures on the Sun's surface, seen in the H-alpha wavelength on August 22, 2003. (Swedish 1-m Solar Telescope (SST) operated by the Royal Swedish Academy of Sciences, Oddbjorn Engvold, Jun Elin Wiik, Luc Roupe van der Voort) #

Zeeman Effect




Breaks "degeneracy" in energy of states with different "spins"


Strength of the field can be measured by the degree of splitting




1.1 hours



about 25,000 km across




11 year cycle




The sun actually changes its luminosity a little ...

Solar Cycle Variations



Variability in solar irradiance was undetectable prior to satellite observations starting in 1978

Sunspot activity as a function of latitude




<http://science.msfc.nasa.gov/sohl/pad/pad/solar/images/bfiv.tif>

NASA/NSSTC/HATHAWAY 2004.02

Spots start to happen at high and low latitude and then migrate towards the equator


Radial magnetic field



Hathaway/NASA/MSFC 2009/09


Complete cycle takes 22 years and the sun's magnetic field reverses every 11 years

Periodic variation in sunspot number



Cause - Multipolar field variation (quadrupole instead of dipole)?
Fluctuations at tachocline?

Little Ice Age



The exact cause of the solar cycle is not well understood, but it is known that the magnetic field of the sun (or at least its surface field) goes through reversals every ~ 11 years. The whole cycle takes 22 years.

In the "Babcock model", the cycle is caused by the *differential rotation of the sun*. In three years the equatorial regions go round 5 additional revolutions compared with the polar ones. This winds up the field and creates stress that is released in part by surface activity (flares, sunspots, etc).


Rotation:

26.8 d at equator

31.8 d at 75° latitude


Rotation:
 26.8 d at equator
 31.8 d at 75° latitude

This differential rotation exists only in the convection zone. The radiative core rotates rigidly.



Babcock Model


Magnetic Field



Babcock Model

Magnetic Field


Differential Rotation



Babcock Model

Magnetic Field





Differential Rotation





Babcock Model

Magnetic
Field

Differential
Rotation




radiation



The smallest loop is 3 times the size of the Earth

Solar Flares




Energy stored in magnetic field

$$E = B^2 / 8\pi \text{ erg cm}^{-3}$$


with B in Gauss

Can be released by "reconnection"




http://sprg.ssl.berkeley.edu/~tohban/nuggets/?page=article&article_id=14

Coronal Mass Ejection



Chromosphere





The chromosphere is a narrow layer above the photosphere. It has low density $\sim 0.01\%$ that of the photosphere.

Initially the temperature and density both decline above the photosphere, but once 3000 K is reached the temperature rises again to 35,000 K. It is heated by magnetic reconnection.

The chromosphere is a transition region from the photosphere to the corona.


Chromosphere	Corona
~Hydrostatic equilibrium	Dynamic
Helium neutral	Helium ionized
Radiates energy effectively	Inefficient radiator
Emission lines	Emission lines
~10,000 K	~1,000,000 K
Reddish due to H α	Highly ionized, uv, x-rays



Coronal Loop

Picture at 141 Angstroms

Temperature About 10⁶ K




Temperature

Center	15.7 million K
Photosphere	5800 K
Sunspot (umbra)	4240 K
Penumbra	5680 K
Chromosphere	5000 – 20000 K
Corona	0.5 to 3 million K

Density


Mean density entire sun	1.41 g cm ⁻³
Central density	150 g cm ⁻³
Photosphere	10 ⁻⁹ g cm ⁻³
Chromosphere	10 ⁻¹² g cm ⁻³
Corona	10 ⁻¹⁶ g cm ⁻³
Air (earth)	10 ⁻³ g cm ⁻³

Mass loss rate $\sim 10^{-13}$ solar masses/yr



http://www.boston.com/bigpicture/2008/10/the_sun.html
<http://stereo.gsfc.nasa.gov>

Solar wind colliding with a comet April 4, 2007. Captured by NASA satellite STEREO (one of 2)



Effects on earth of solar cycle


- Radio communications and satellite health
- Ozone production and hence uv flux at earth
- Cosmic ray flux
- Aurorae



Carrington Event 1859


https://en.wikipedia.org/wiki/Solar_storm_of_1859

- Aurorae worldwide – as far south as Hawaii
- Telegraph disruptions. Sparks flying. Able to send messages without batteries connected
- Followed observation of a huge solar flare
- Estimated damage today 0.6 to 2.6 trillion dollars
- Similar event on July 23, 2012. Just missed the earth



The last complete reversal was 980,000 yrs ago, but there was almost a reversal 200,000 years

Names are the rock strata where the field is measured



<http://www.astronomycafe.net/qadir/q816.html>

The current magnetic field strength at the Earth's surface of 0.6 Gauss. But long term observations show that it is DECREASING at a rate of about 0.07 percent PER YEAR. This means that in 1500 years from now, it will only be about 35 percent as strong as it is today, and in 4000 years it will have a strength of practically zero.



Magnetic north is currently in northern Canada moving at 10 to 50 km/yr.

In a few decades it will reach Siberia


Solar flares vs solar prominences (the latter are bigger)

<http://science.howstuffworks.com/sun5.htm>

http://science.nasa.gov/science-news/science-at-nasa/2003/29dec_magneticfield/

The Solar Neutrino "Problem"

Hydrogen Burning on the Main Sequence





- pp1 85%
- pp2 15%
- pp3 0.02%

$T_{\text{central}} = 15.7 \text{ Million K}$


Hydrogen Burning on the Main Sequence

In all cases $4p \rightarrow {}^4\text{He} + 2\nu_e + 2e^+$



pp1



pp2



pp3



Neutrino Energies

	Species	Average energy	Maximum energy	
pp1	p+p	0.267 MeV	0.420 MeV	
pp2 and pp3	${}^7\text{Be}$	0.383 MeV	0.383 MeV	10%
		0.861	0.861	90%
pp3	${}^8\text{B}$	6.735 MeV	15 MeV	

In the case of ${}^8\text{B}$ and p+p, the energy is shared with a positron hence there is a spread. For ${}^7\text{Be}$ the electron capture goes to two particular states in ${}^7\text{Li}$ and the neutrino has only two energies

CAN WE DETECT THESE NEUTRINOS?

Since 1965, experiments have operated to search for and study the neutrinos produced by the sun - in order to:


- Test solar models
- Determine the central temperature of the sun

The flux of neutrinos from ^8B is sensitive to T^{18}


- Learn new particle physics

DETECTORS

The chlorine experiment – Ray Davis – 1965 - ~1999




i.e., a neutron inside of ^{37}Cl is turned into a proton by a weak interaction involving an incident neutrino




DETECTORS

The chlorine experiment – Ray Davis – 1965 - ~1999



The gallium experiments (GALLEX and SAGE) – 1991 – 1997 and 1990 – 2001



Homestake Gold Mine
Lead, South Dakota

4850 feet down

tank 20 x 48 feet
615 tons (3.8×10^5 liters)
 C_2Cl_4


Threshold 0.814 MeV

Half-life ^{37}Ar = 35.0 days

Neutrino sensitivity
 ^7Be , ^8B

8×10^{30} atoms of Cl

Nobel Prize 2002




Calibrated using radioactive ^{51}Cr neutrino source


In Gran Sasso Tunnel – Italy
 3300 m water equivalent
 30.3 tons of gallium in GaCl_3 -
 HCl solution
 $^{71}\text{Ga} + \nu_e \rightarrow ^{71}\text{Ge} + e^-$
 Threshold 0.233 MeV
 Sees pp, ^7Be , and ^8B .

DETECTORS


The chlorine experiment – Ray Davis – 1965 - ~1999



The gallium experiments (GALLEX and SAGE) –
 1991 – 1997 and 1990 – 2001




Kamiokande II - 1996 – 2001




Inelastic scattering of neutrinos on electrons in water. Threshold 9 MeV. Scattered electron emits characteristic radiation.


Kamiokande II (in Japanese Alps) 1996 - 2001




Depth 1 km
 Detector H_2O
 Threshold 9 MeV
 Sensitive to ^8B
 20' photomultiplier
 tubes
 Measure Cerenkov
 light
 2.3×10^{32} electrons



Super-Kamiokande (Japan)
 50,000 tons of water
 11,146 20" light detectors
 0.6 mi down




The Sun - 1999
 (First picture in neutrinos)



This "picture" was taken using data from the Kamiokande 2 neutrino observatory. It contains data from 504 nights (and days) of observation. The observatory is about a mile underground.

Each pixel is about a degree and the whole frame is 90° x 90°.

Sudbury Neutrino Observatory



6800 ft down
 1000 tons D₂O.
 20 m diameter
 Sudbury, Canada
 Threshold 5 MeV
 Sees ⁸B decay but can see all three kinds of neutrinos

$$\nu_e, \nu_\mu, \nu_\tau$$



Particle physics aside:

Three Generations of Matter (Fermions)

	I	II	III	
mass →	2.4 MeV	1.27 GeV	171.2 GeV	0
charge →	2/3	2/3	2/3	0
spin →	1/2	1/2	1/2	1
name →	u up	c charm	t top	γ photon
	4.8 MeV	104 MeV	4.2 GeV	0
	-1/3	-1/3	-1/3	0
	1/2	1/2	1/2	1
Quarks	d down	s strange	b bottom	g gluon
	<2.2 eV	<0.17 MeV	<15.5 MeV	91.2 GeV
	0	0	0	0
	1/2	1/2	1/2	1
	ν_e electron neutrino	ν_μ muon neutrino	ν_τ tau neutrino	Z weak force
	0.511 MeV	105.7 MeV	1.777 GeV	80.4 GeV
	-1	-1	-1	±1
	1/2	1/2	1/2	1
Leptons	e electron	μ muon	τ tau	W weak force

emitted by pp-cycle
 cosmology limits the sum of the 3 neutrino masses to < 1 eV


Bosons (Forces)





<http://www.sno.phy.queensu.ca/sno/sno2.html> - interactions

Neutrino interactions with heavy water $D_2O = {}^2H_2O$

Electron neutrino



All neutrinos ($e, \mu,$ and τ) with energy above $2.2 \text{ MeV} = BE({}^2H)$



Results from SNO – 2002 (Sudbury turned off in 2006)

The flux of electron flavored neutrinos above 5 MeV (i.e., only pp3 = 8B neutrinos) is

$$1.76 \pm 0.1 \times 10^6 \text{ cm}^{-2} \text{ s}^{-1}$$

But the flux of μ and τ flavored neutrinos is

$$3.41 \pm 0.64 \times 10^6 \text{ cm}^{-2} \text{ s}^{-1}$$

Nobel Prize in Physics - 2002

Standard Solar Model 8B neutrinos

$$5.05^{+1.01}_{-0.81} \times 10^6 \text{ neutrinos cm}^{-2} \text{ s}^{-1}$$

The explanation of the solar neutrino “problem” is apparently *neutrino flavor mixing*.

http://en.wikipedia.org/wiki/Neutrino_oscillation

A flux that starts out as pure electron-”flavored” neutrinos at the middle of the sun ends up at the earth as a mixture of electron, muon, and tauon flavored neutrinos in comparable proportions.

The transformation occurs in the sun and is complete by the time the neutrinos leave the surface. The transformation affects the highest energy neutrinos the most (MSW-mixing).

Such mixing requires that the neutrino have a very small but non-zero rest mass. This is different than in the so called “standard model” where the neutrino is massless. The mass is less than about 10^{-5} times that of the electron. (Also observed in earth’s atmosphere and neutrinos from reactors).

New physics.... (plus we measure the central temperature of the sun very accurately – 15.71 million K)